Table 14.2. TMD Clinician Self-Assessment for Adherence to RCT Study Protocols

University of Washington - Department of Oral Medicine Clinical Examination & RDC for TMD Database

Name: #:	: Date:
TREATMENT & MANAGEMENT PLAN: 1 = New 2	= Continue 3 = Modify Rvt: Avt:
Physical Medicine, Physiotherapy, Muscle and Jaw Relaxation:  Keep jaws relaxed, teeth separated, lips apart & check frequently to assure the muscles are relaxed.  Reduce parafunctional jaw habits. Don't bite fingernails or cheeks, or hold pencils or other items in mouth.  Soft diet avoid crunchy foods avoid chewy foods cut food up small bites avoid opening wide Don't stretch jaw muscles by moving jaw side to side. Don't posture jaw forward or to the side.  Relax jaw muscles by placing tip of tongue behind lower front teeth. Let tongue go completely relaxed (check often).  Open as wide as possible without pain. Hold sec., close halfway & rest for 5 sec. ( repetitions, times/day).  Apply cold pack or heat pack (moist heat preferred) to areas of pain. ( minutes times/day).	
Keep shoulders down and relaxed - check often to see if shoulder muscles are relaxed.  Stretch neck muscles by tipping head down slowly until chin touches chest (repeat times). Keep shoulders down.  With chin to chest, tip head side to side & hold for 15 seconds. ( repetitions , times /day).  Apply	
General and Behavioral Management:  Identify stressors and make a plan to manage them.  Obtain and read a book on stress management.  Improve sleep pattern & amount of sleep.  Massage the areas of pain daily by hand or machine.	Exercise 30 min., 5 times/wk., walking or aerobic.  Avoid caffeine to enhance relaxation.  Relaxation  Practice expressive writing.  Develop a pain management personal plan.
Medications and dosages:  NSAIDS: Narcotic #1: Narcotic #2: Mus. Relaxant: Anxiolytic:	Anti.Depress: Sedative: Other #1: Other #2: Change other medication:
Special Procedures:  Active jaw opening exercises:  Spray and stretch:  Joint manipulation:  Spot occlusal adjustment:	Trigger point injection:  Nerve block:  Joint inject:  Other:
Splints and Appliances:         Splint pre-authorization       Splint impression         Seat hard splint       Seat soft splint         Adjust existing splint       Fabricate special appliance	Provide soft splint blank Use splint hrs. while awake, hrs. at night Other:

Signature: